

Problem E - Extreme Divisors

Let us define the functions $d(n)$ and $\sigma(n)$ as

$d(n)$ = number of divisors of n

$\sigma(n)$ = summation of divisors of n

Here divisors of n include both 1 and n . For example divisors of 6 are 1, 2, 3 and 6. So $d(6) = 4$ and $\sigma(6) = 12$

Now let us define two more function $g(a,b,k)$ and $h(a,b,k)$ as

$$g(a,b,k) = \sum_{i} d(i)$$

$$h(a,b,k) = \sum_{i} \sigma(i)$$

Where $a \leq i \leq b$ and i is divisible by k .

For example, $g(5,12,3) = d(6) + d(9) + d(12) = 4 + 3 + 6 = 13$ and $h(5,12,3) = \sigma(6) + \sigma(9) + \sigma(12) = 12 + 13 + 28 = 53$. Given a, b, k you have to calculate $g(a,b,k)$ and $h(a,b,k)$.

Input

The first line of the input file contains an integer T ($T \leq 75$) which denotes the total number of test cases. The description of each test case is given below:

Three integers in a line. First integer is contains a , second integer is b and third integer is k . You may assume $0 < a \leq b \leq 10^{12}$, $0 < k < 2000$.

Output

For each test case print one line containing $g(a,b,k)$ and $h(a,b,k)$ separated by a space as defined above. As output may be very large print the output modulo 2^{64} .

Sample Input

```
2
5 12 3
1 100 3
```

Sample Output

```
13 53
217 3323
```

Problem Setter: Tanaeem M Moosa
Special Thanks: Shakil Ahmed
Next Generation Contest 6