


## Problem G

# Go up the Ultras

The topographic prominence of a peak is a measure of special interest to mountain climbers and can be defined as follows: the prominence of a peak  $p$  with altitude  $h$ , relative to the sea level, is the greatest  $d$  such that any path on the terrain from  $p$  to any strictly higher peak will pass through a point of altitude  $h - d$ . If there is no strictly higher peak, then the prominence is  $h$  itself. Those peaks with topographic prominence greater than or equal to 150000 centimeters (precision is of great importance to climbers!) have a special name: they are called “Ultras”.

You have to write a program that identifies all the Ultras that occur in a two dimensional profile of a mountain range represented as a sequence of points. Note that the horizontal distance between points is not important; all that you need is the altitude of each point. In the picture below, the Ultras are the points 7, 12, 14, 20 and 23.



### Input

The first line contains an integer  $N$  ( $3 \leq N \leq 10^5$ ) representing the number of points in the profile. The second line contains  $N$  integers  $H_i$  indicating the altitudes (in centimeters) of the points, in the order in which they appear in the profile ( $0 \leq H_i \leq 10^6$  for  $i = 1, 2, \dots, N$ ). Consecutive points have different altitudes ( $H_i \neq H_{i+1}$  for  $i = 1, 2, \dots, N - 1$ ), while the first and the last points are at sea level ( $H_1 = H_N = 0$ ). You may assume that the profile contains at least one Ultra.

### Output

Output a line with the indices of all the Ultras in the mountain range, in the order in which they appear in the profile.

<p><b>Sample input 1</b></p> <p>5</p> <p>0 10000 100000 884813 0</p>	<p><b>Sample output 1</b></p> <p>4</p>
<p><b>Sample input 2</b></p> <p>7</p> <p>0 100000 0 200000 180000 200000 0</p>	<p><b>Sample output 2</b></p> <p>4 6</p>